



FLWEMS Paramedic Medication Information For:

ASPRIN

(Endodan)

(ASS-pih-rin)

Endodan Oxycodan★ Percodan Percodan-Demi★ (C-II) (Rx)

Classification: Analgesic

See Also

See also *Acetylsalicylic acid* and *Narcotics Analgesics*.

Content

Each tablet contains: *Narcotic analgesics*: oxycodone, 4.5 mg and oxycodone terephthalate, 0.38 mg.
Analgesic: Aspirin, 325 mg.

Uses

Relief of moderate to moderately severe pain.

Contraindications

Use in children and during pregnancy.

Special Concerns

Oxycodone can produce drug dependence and has abuse potential. The respiratory depressant effects of oxycodone can be exaggerated in clients with head injury, other intracranial lesions, or a preexisting increase in intracranial pressure. Use with caution in clients who are elderly, are debilitated, have severely impaired hepatic or renal function, are hyperthyroid, have Addison's disease, have prostatic hypertrophy, or have urethral stricture. Use for acute abdominal conditions may obscure the diagnosis or clinical course.

Side Effects

Commonly, dizziness, lightheadedness, N&er:V, and sedation; these effects are more common in ambulatory clients than in nonambulatory clients. Other side effects include euphoria, dysphoria, constipation, skin rash, and pruritus. See also individual components.

Drug Interactions

Aspirin / ↑ Effect of anticoagulants *CNS depressants* (including other narcotic analgesics, phenothiazines, antianxiety drugs, sedative-hypnotics, anesthetics, alcohol) / Additive CNS depression

Dosage

•Tablets *Analgesic*.

Adults: 1 tablet q 6 hr as needed for pain; higher doses may be required in cases of more severe pain or in those who have become tolerant to the analgesic effect of narcotics.

END OF INFORMATION – NOTHING FOLLOWS